



Length-Weight and Length-Length Relationships of the *Scorpaena* Species (Actinopteri: Scorpaenidae) in Izmir Bay (Aegean Sea of Turkey)

Bahar Bayhan¹ • Burcu Taylan¹ • Oğulcan Heral¹

¹ Ege University, Faculty of Fisheries, Department of Basic Science, Bornova, İzmir, Turkey, bahar.bayhan@ege.edu.tr; burcu.taylan@ege.edu.tr; ogulcan.heral@gmail.com

✉ Corresponding Author: burcu.taylan@ege.edu.tr

Please cite this paper as follows:

Bayhan, B., Taylan, B., & Heral, O. (2022). Length-Weight and Length-Length Relationships of the *Scorpaena* Species (Actinopteri: Scorpaenidae) in Izmir Bay (Aegean Sea of Turkey). *Acta Natura et Scientia*, 3(1), 51-58. <https://doi.org/10.29329/actanatsci.2022.351.06>

ARTICLE INFO

Article History

Received: 27.04.2022

Revised: 11.05.2022

Accepted: 12.05.2022

Available online: 14.06.2022

Keywords:

Scorpaena

Length-length relationship

Length-weight relationship

Aegean Sea

Izmir Bay

ABSTRACT

In this study, 173 specimens of *Scorpaena porcus* (n=50), *Scorpaena elongata* (n=20), *Scorpaena scrofa* (n=101), *Scorpaena notata* (n=2) were obtained from commercial fishermen in the Aegean Sea between 2017 and 2019. The length-weight relationships were determined for *S. porcus*, *S. scrofa* and *S. elongata* as $W=0.0119TL^{3.182}$, $W=0.0176TL^{3.004}$ and $W=0.0656TL^{2.579}$, respectively. Negative allometry was found for *S. scrofa* and *S. elongate*, whereas positive allometry was present for *S. porcus*. However, with less than 5 individuals *S. notata* showed maximum, minimum, and mean total length and weight values. *S. notata* (2 individuals) had a mean total length and weight of 15.7-16.6 cm and 72.31-93.26 g. The results also demonstrated that the length-length relationships were strongly correlated for all *Scorpaena* species ($r^2>0.980$, $P<0.05$).

INTRODUCTION

Aquatic biologists and fisheries managers have frequently used length-weight relationships (LWRs) to evaluate fish stocks, the ecology of fisheries and the population dynamics in aquatic ecosystems, as well as in aquatic environments that are semi-controlled (Ricker, 1968). Length-weight relationships help appraise the general health, reproduction history, life cycle and condition of fish species (Pauly, 1993). They can be helpful in comparing local and interregional morphologies in both populations and species. In addition, length-weight models can be used to assess the ecological health and productivity levels of aquatic

ecosystems (Deekae & Abowei, 2010). Models such as these serve to assess the health and growth patterns of fish (Muchlisin et al., 2010; Ndiaye et al., 2015). There are 231 species of Scorpaenidae worldwide (Fricke et al., 2022), while there are eight species found off the Turkish coasts (Karataş et al., 2021).

It is often possible in fisheries studies to measure fish length more easily and quickly than to measure mass. Having a knowledge of the length-weight relationship facilitates the determination of the mass if only the length is known. When working in the field, tail flukes are often cut; this can make it difficult to obtain accurate measurements of total length.

Knowing the length enables the researcher to work out total length.

The length-weight relationships of the different *Scorpaena* fish populations have been indicated in previous studies in the seas around Turkey (Koca, 2002; Karakulak et al., 2006; Özeydin & Taşkavak, 2006; Ismen et al., 2007; Demirhan & Can, 2007; Alpaslan et al., 2007; Ak et al., 2009; Keskin & Gaygusuz, 2010; Akalın et al., 2011; Bok et al., 2011; Kasapoğlu & Düzgünleş, 2013; Bilge et al., 2014; Deval et al., 2014; Altın et al., 2015) and other regions (Jardas & Pallaoro, 1992; Petrakis & Stergiou, 1995; Dulčić & Kraljevic, 1996; Stergiou & Moutopoulos, 2001; Moutopoulos & Stergiou, 2002; Mendes et al., 2004; Ferreira et al., 2008; Crechriou et al., 2013; Moutopoulos et al., 2013; Dimitriadis & Konstantinidou, 2018; Meiners-Mandujano et al., 2018; Miled-Fathali et al., 2019).

Nevertheless, the length-weight parameters of the same species can differ in the population as a result of fishing, reproductive activities and feeding etc. It is thus necessary to be aware of the length-weight relationships of fish caught in a specific place during a specific period of time.

The aim of the current study was to determine the length-weight and length-length relationships of *Scorpaena* species caught in Izmir Bay in 2017 and 2019.

MATERIAL AND METHODS

In this study, 173 specimens of *S. porcus* (n=50), *S. elongata* (n=20), *S. scrofa* (n=101), *S. notata* (n=2) were obtained from commercial fishermen in the Aegean Sea between 2017 and 2019. Total length (TL) and standard length (SL) of each specimen was measured

using a 0.1 cm class interval. The total weight (W) of each individual was measured to the nearest 0.01 g. The total length of all specimens was employed to work out the length-weight relationship (LWR). This was calculated by log transformed data log (Eq. 1):

$$W = \log a + b \log L \tag{1}$$

where, W is the total weight (g), L is the total length (cm), a is the intercept and b is the slope or allometric coefficient (Ricker, 1979). The statistical significance level of r² was predicted using linear regressions on the transformed equation. An allometric coefficient (b) larger or smaller than 3.0 shows an allometric growth (Bagenal & Tesch, 1978). Value b is > 3, which can be said to show a positive allometric growth. However, b<3 demonstrated a negative allometric growth or isometric growth when it was equal to 3.0. Moreover, the SL vs TL relationships were calculated using linear regressions. Length-length relationship (LLR) was measured as Eq. (2) to standardize the length values used in other studies.

$$TL = a + bSL \tag{2}$$

Covariance analysis was conducted to test for probable significant differences in both slope and intercept. Statistical analyses were evaluated at a significance level of P<0.05

RESULTS

The study examined 173 scorpaenid species of fish with 101 *Scorpaena scrofa*, 50 *Scorpaena porcus*, 20 *Scorpaena elongata* and 2 *Scorpaena notata*.

Table 1. Descriptive statistics and estimated parameters of length-weight relationships for four *Scorpaena* species in Izmir Bay (Aegean Sea) from 2017 to 2019 (N: number of individuals, a: intercept, b: slope, r²: coefficient of determination)

Species	N	TL range (cm)	W range (g)	a	B	95% C.I. of b	r ²	Growth type
<i>S. scrofa</i>	101	17.3-33.1	95.37-708.91	0.0176	3.004	2.892-3.116	0.963	-allometry
<i>S. porcus</i>	50	12.6-28.4	33.76-516.03	0.0119	3.182	3.042-3.322	0.972	+allometry
<i>S. elongata</i>	20	19.7-28.3	134.57-358.68	0.0656	2.579	2.347-2.811	0.958	-allometry
<i>S. notata</i>	2	15.7-16.6	72.31-93.26	-	-	-	-	-




Figure 1. Length-weight relationships for three Scorpaena species from Izmir Bay (Central Aegean Sea, Turkey).

Table 2: Length-length relationships between total length (TL), and standard length (SL) of *Scorpaena* species in Izmir Bay (Aegean Sea) from 2017 to 2019 (N: number of individuals, *a*: intercept, *b*: slope, *r*²: coefficient of determination)

Species	Equation	N	<i>a</i>	<i>b</i>	<i>r</i> ²
<i>S. scrofa</i>	TL=a+bSL	101	0.3354	0.7744	0.986
<i>S. porcus</i>	TL=a+bSL	50	0.6342	0.8171	0.989
<i>S. elongata</i>	TL=a+bSL	20	0.3588	0.7827	0.990

Table 3. LWRs of three *Scorpaena* species from different areas (N: number of individuals, *a*: intercept, *b*: slope, *r*²: coefficient of determination)

Species	References	Country, Locality	N	Length range (TL, cm)	<i>a</i>	<i>b</i>	<i>r</i> ²
<i>S. porcus</i>	Jardas & Pallaoro (1992)	Croatia, Adriatic	-	5.8-30.0	0.0171	3.034	0.995
	Petrakis & Stergiou (1995)	Greece, Euboikos Gulf	100	7.0-23.5	0.0291	2.840	0.970
	Stergiou & Moutopoulos (2001)	Greece, Aegean Sea	231	11.5-40.5	0.0236	2.887	0.950
	Stergiou & Moutopoulos (2001)	Greece, Evvoikos	100	7.0-23.5	0.0290	2.840	0.970
	Stergiou & Moutopoulos (2001)	Greece, Cyclades	231	11.5-40.5	0.0236	2.887	0.951
	Koca (2002)	Turkey, Sinop Peninsula	633	12.0-25.0	0.0540	2.590	0.931
	Morey et al. (2003)	Spain, Coast of Iber	-	6.1-35.5	0.0183	3.020	0.969
	Karakulak et al. (2006)	Turkey, Gökçeada Island	255	8.0-27.3	0.0250	2.915	0.931
	Özaydin & Taskavak (2006)	Turkey, İzmir Bay	50	14.1-25.6	0.0210	3.004	0.960
	Ismen et al. (2007)	Turkey, Saroz Bay	-	10.0-22.0	0.0256	2.878	0.993
	Demirhan & Can (2007)	Turkey, Black Sea	470	4.6-17.5	0.0124	3.190	0.940
	Alpaslan et al. (2007)	Turkey, Dardanelles	-	10.5-32.0	0.0230	2.960	0.700
	Ak et al. (2009)	Turkey, Coast of Trabzon	351	5.0-34.2	0.0090	3.272	0.880
	La Mesa et al. (2010)	Italy, Adriatic artificial reef	210	8.0-27.0	0.0202	3.030	0.950
	La Mesa et al. (2010)	Italy, Adriatic nature reef	134	11.0-22.1	0.0257	2.940	0.950
	La Mesa et al. (2010)	Italy, Adriatic offshore	71	13.5-28.0	0.0136	3.120	0.970
	Keskin & Gaygusuz (2010)	Turkey, Erdek Bay	45	4.9-19.0	0.0158	3.088	0.983
	Akalın et al. (2011)	Turkey, İzmir Bay	221	7.0-27.0	0.0209	2.987	0.993
	Bok et al. (2011)	Turkey, Sea of Marmara	15	17.3-21.4	0.0067	3.343	0.944
	Crec'hriou et al. (2013)	France, Coast of Catalan	111	9.5-32.0	0.0260	2.920	0.968
Moutopoulos et al. (2013)	Greece, Korinthiakos Bay	103	10.5-31.8	0.0192	2.965	0.969	
Kasapoglu & Duzgunes (2013)	Turkey, Black Sea	136	8.5-29.2	0.0173	3.034	0.980	
Bilge et al. (2014)	Turkey, South Aegean Sea	63	7.7-26.8	0.0170	3.045	0.986	
Miled-Fathali et al. (2019)	Tunisian, Tunisian Bay	70	8.6-23.6	0.0295	2.830	0.980	
This study	Turkey, İzmir Bay	50	12.6-28.4	0.0119	3.182	0.972	
<i>S. scrofa</i>	Dulčić & Kraljevic (1996)	Croatia, Eastern Adriatic	125	19.7-53.6	0.0156	3.298	0.960
	Stergiou & Moutopoulos (2001)	Greece, Cyclades	37	10.7-32.2	0.0169	2.999	0.980
	Moutopoulos & Stergiou (2002)	Greece, Aegean Sea	37	10.7-32.2	0.0169	2.999	0.980
	Mendes et al. (2004)	Portugal, from Nazaré to St André	22	14.3-42.8	0.0121	3.124	0.933
	Karakulak et al. (2006)	Turkey, Gökçeada Island	15	12.3-39.1	0.0180	3.005	0.985
	Özaydin & Taskavak (2006)	Turkey, İzmir Bay	129	8.2-30.1	0.0291	2.796	0.990
	Ferreira et al. (2008)	Madeira islands, Atlantic	12	20.0-46.0	0.0153	3.039	0.974
	Crec'hriou et al. (2013)	France, Coast of Catalan	32	14.0-49.0	0.0260	2.890	0.993
	Moutopoulos et al. (2013)	Greece, Korinthiakos Bay	53	11.7-36.8	0.0169	3.002	0.981
	Bilge et al. (2014)	Turkey, Southern Aegean Sea	74	9.6-42.9	0.0217	2.964	0.996
	Altın et al. (2015)	Turkey, Gökçeada Island	16	6.3-13.2	0.0120	3.135	0.983
	Dimitriadis & Konstantinidou (2018)	Greece, Zakintos Island	335	10.5-39.7	0.0260	2.867	0.980
	Miled-Fathali et al. (2019)	Tunisian, Tunisian Bay	61	11.3-38.3	0.0205	2.947	0.950
	This study	Turkey, İzmir Bay	50	17.3-33.1	0.0176	3.004	0.963
<i>S. elongata</i>	Deval et al. (2014)	Turkey, Antalya Bay	114	5.6-42.1	0.0230	2.880	0.996
	Meiners-Mandujano et al. (2018)	Morocco	45	18.6-41.2	0.0148	3.017	0.971
	Miled-Fathali et al. (2019)	Tunisian, Tunisian Bay	22	14.2-50.0	0.0127	3.077	0.990
	This study	Turkey, İzmir Bay	20	19.7-28.3	0.0656	2.579	0.958

All the specimens of *Scorpaena scrofa* examined showed length group values of 28-94 cm with 17.3-33.1 cm length. The mean total length and weight values of the species were 22.32 ± 0.32 cm and 269.93 ± 11.21 g, respectively. The total length-weight relationship was found to be $W = 0.0176TL^{3.004}$ ($r^2 = 0.963$) with a negative allometric growth. 50 individuals of *S. porcus* species exhibited minimum maximum length range of 12.6-28.4 cm. Its mean total length and weight values were 19.29 ± 0.52 cm and 164.88 ± 15.10 g, respectively. The total length and weight relationship was $W = 0.0119TL^{3.182}$ ($r^2 = 0.972$) with a positive growth. 20 individuals of *S. elongata* had a minimum maximum length range of 19.7-28.3 cm. Its mean total length and weight values were 24.61 ± 0.56 cm and 258.73 ± 14.87 g, respectively. Total length weight relationship was $W = 0.0656TL^{2.579}$ ($r^2 = 0.958$) with a positive allometric growth. Nevertheless, with fewer than five individuals *S. notata* showed minimum, maximum and mean total length and weight values. *S. notata* (two individuals) had mean total length and weight of 15.7-16.6 cm and 72.31-93.26 g, respectively (Table 1, Figure 1). Of the LLRs shown in Table 2, all were highly significant ($P < 0.05$), and all coefficient of determination values were larger than 0.980.

DISCUSSION

The current study examined 173 individuals from three *Scorpaena* species. Previous studies on the length-weight relationship for the three species examined in the current study are presented in Table 3.

The b values in LWRs lie between 2.5 and 3.5 (Froese, 2006) or from 2 to 4 (Tesch, 1971). In the current study, the b values of the fish species studied fell within these ranges, as expected. The b value for *S. porcus* ranged from 2.590 to 3.190 in other areas, while the value ranged from 2.796 to 3.135 for *S. scrofa*. The b value for *S. elongata* was determined to be 2.579. Although this value is different from those in the studies of Meiners-Mandujano et al. (2018) and Miled-Fathali et al. (2019), it is similar to Deval et al. (2014) (Table 3).

Length-weight relationships may vary in different seasons, and they are influenced by factors such as temperature, size, maturity, food availability and

salinity. Differences in b values can be put down to one or a mixture of these factors, including the number of specimens studied, the effects of the region and/or season, and differences in the ranges of lengths of the specimens; the length of time in which the sample was collected can also be added to these (Moutopoulos & Stergiou, 2002). Furthermore, the extent of sexual maturity, fullness or emptiness of the gut, diet, sampling techniques, and the number of specimens and period of sampling can also have an effect on its value (Wootton, 2003; Eagderi et al., 2020).

CONCLUSION

To conclude, this study has given new information regarding LWRs and LLRs for three *Scorpaena* species from Izmir Bay (the Central Aegean Sea of Turkey). It has provided additional data concerning these previously reported species, and will thus be of use in future biological studies.

ACKNOWLEDGEMENTS

The study was carried out with the financial support of the Ege University Scientific Research Project (2017/SUF/15), Izmir, Turkey.

Compliance with Ethical Standards

Authors' Contributions

Conceptualization: BB, BT. Data collection and analysis: BT. Conducting the laboratory experiments: BT, OH. Writing: BB, BT.

Conflict of Interest

The authors declare that there is no conflict of interest.

Ethical Approval

For this type of study, formal consent is not required.

REFERENCES

- Ak, O., Kutlu, S., & Aydin, I. (2009). Length-weight relationship for 16 fish species from the Eastern Black Sea, Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*, 9, 125-126.

- Akalın, S., İlhan, D., Ünlüoğlu, A., Tosunoğlu, Z., & Özeydin, O. (2011). İzmir Körfezi'nde iki iskorbüt türünün (*Scorpaena notata* ve *Scorpaena porcus*) boy-ağırlık ilişkisi ve metrik-meristik özellikleri [Length-weight relationship and metric-meristic characteristics of two scorpion fishes (*Scorpaena notata* and *Scorpaena porcus*) in İzmir Bay]. *Journal of Fisheries Science.com*, 5(4), 291-299. <https://doi.org/10.3153/jfscom.2011033>
- Alpaslan, M., Tekinay, A. A., & Çınar, Y. (2007). Çanakkale Boğazi'nda dağılım gösteren iskorbüt balığı (*Scorpaena porcus* Linnaeus, 1758)'nin bazı büyüme parametreleri [Some growth parameters of scorpion fish (*Scorpaena porcus* Linnaeus, 1758) in Dardanelles]. *E.Ü. Su Ürünleri Dergisi*, 24(3-4), 219-223.
- Altun, A., Ayyıldız, H., Kale, S., & Alver, C. (2015). Length-weight relationships of forty-nine fish species from shallow waters of Gökçeada Island, northern Aegean Sea. *Turkish Journal of Zoology*, 39(5), 971-975. <https://doi.org/10.3906/zoo-1412-15>
- Bagenal, T. B., & Tesch, F. W. (1978). Age and growth. In T. Bagenal (Ed.), *Methods for assessment of fish production in fresh waters* (pp. 101-136). (3rd ed.). Blackwell Science Publications.
- Bilge, G., Yapıcı, S., Filiz, H., & Cerim, H. (2014). Weight-length relations for 103 fish species from the southern Aegean Sea, Turkey. *Acta Ichthyologica Piscatoria*, 44(3), 263-269. <https://doi.org/10.3750/AIP2014.44.3.11>
- Bok, T. D., Gokturk, S. D., Kahraman, A. E., Alicli, T. Z., Acun, T., & Ates, C. (2011). Length-weight relationships of 34 fish species from the Sea of Marmara, Turkey. *Journal of Animal and Veterinary Advances*, 10(23), 3037-3042. <https://doi.org/10.3923/javaa.2011.3037.3042>
- Crechriou, R., Neveu, R., & Lenfant, P. (2013). Length-weight relationship of main commercial fishes from the French Catalan coast. *Journal of Applied Ichthyology*, 29(5), 1191-1192. <https://doi.org/10.1111/jai.12320>
- Deekae, S. N., & Abowei, J. F. N. (2010). *Macrobrachium macrobrachion* (Herklots, 1851) length-weight relationship and Fulton's condition factor in Luubara creek, Ogoni land, Niger Delta, Nigeria. *International Journal of Animal and Veterinary Advances*, 2(4), 155-162.
- Demirhan, S. A., & Can, M. F. (2007). Length-weight relationships for seven fish species from the southeastern Black Sea. *Journal of Applied Ichthyology*, 23(3), 282-283. <https://doi.org/10.1111/j.1439-0426.2007.00835.x>
- Deval, M. C., Güven, O., Saygu, I., & Kabapçıoğlu, T. (2014). Length-weight relationships of 10 fish species found off Antalya Bay, eastern Mediterranean. *Journal of Applied Ichthyology*, 30(3), 567-568. <https://doi.org/10.1111/jai.12382>
- Dimitriadis, C., & Fournari-Konstantinidou, I. (2018). Length-weight relations for 20 fish species (Actinopterygii) from the Southern Ionian Sea, Greece. *Acta Ichthyologica et Piscatoria*, 48(4), 415-417. <https://doi.org/10.3750/AIEP/02466>
- Dulčić, J., & Kraljevic, M. (1996). Weight-length relationship for 40 fish species in the eastern Adriatic (Croatian waters). *Fisheries Research*, 28(3), 243-251. [https://doi.org/10.1016/0165-7836\(96\)00513-9](https://doi.org/10.1016/0165-7836(96)00513-9)
- Eagderi, S., Mouludi-Saleh, A., & Cicek, E. (2020). Length-weight relationship of ten species of Leuciscinae sub-family (Cyprinidae) from Iranian inland waters. *International Aquatic Research*, 12(2), 133-136. [https://doi.org/10.22034/iar\(20\).2020.1891648.1004](https://doi.org/10.22034/iar(20).2020.1891648.1004)
- Ferreira, S., Sousa, R., Delgado, J., Carvalho, D., & Chada, T. (2008). Weight-length relationships for demersal fish species caught off the Madeira archipelago (eastern-central Atlantic). *Journal of Applied Ichthyology*, 24(1), 93-95. <https://doi.org/10.1111/j.1439-0426.2007.01027.x>
- Fricke, R., Eschmeyer, W. N., & Fong, J. D. (2022). Eschmeyer's Catalog of Fishes. Species by family/subfamily. Retrieved on March 21, 2022, <https://researcharchive.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp>
- Froese, R. (2006). Cube law, condition factor and weight-length relationships: History, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22(4), 241-253. <https://doi.org/10.1111/j.1439-0426.2006.00805.x>

- Ismen, A., Ozen, O., Altinagaç, U., Ozekinci, U., & Ayaz, A. (2007). Weight-length relationships of 63 fish species in Saros Bay, Turkey. *Journal of Applied Ichthyology*, 23(6), 707-708. <https://doi.org/10.1111/j.1439-0426.2007.00872.x>
- Jardas, I., & Pallaoro, A. (1992). Age and growth of black scorpionfish *Scorpaena porcus* L. 1758 in the Adriatic Sea. *Rapports et Procès-Verbaux des Réunions de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée*, 33, 296.
- Karakulak, F. S., Erk, H., & Bilgin, B. (2006). Length-weight relationships for 47 coastal fish species from the northern Aegean Sea, Turkey. *Journal of Applied Ichthyology*, 22(4), 274-278. <https://doi.org/10.1111/j.1439-0426.2006.00736.x>
- Karataş, A., Filiz, H., Erciyas-Yavuz, K., Saniye Cevher Özeren, S., & Tok, C. V. (2021). The Vertebrate biodiversity of Turkey. In M. Öztürk, V. Altay, & R. Efe (Eds.), *Biodiversity, conservation and sustainability in Asia* (pp. 175-274). Springer Nature Switzerland. https://doi.org/10.1007/978-3-030-59928-7_10
- Kasapoglu, N., & Duzgunes, E. (2013). Length-weight relationships of marine species caught by five gears from the Black Sea. *Mediterranean Marine Science*, 15(1), 95-100. <https://doi.org/10.12681/mms.463>
- Keskin, Ç., & Gaygusuz, Ö. (2010). Length-weight relationships of fishes in shallow waters of Erdek Bay (Sea of Marmara, Turkey). *IUFS Journal of Biology*, 69(2), 87-94.
- Koca, H. U. (2002). A study on the determination of some parameters of the scorpion fish (*Scorpaena porcus* Linne., 1758) caught by bottom nets in the area of Sinop in terms of fishery biology. *Turkish Journal of Veterinary and Animal Sciences*, 26, 65-69.
- La Mesa, M., Scarcella, G., Grati, F., & Fabi, G. (2010). Age and growth of the black scorpionfish, *Scorpaena porcus* (Pisces: Scorpaenidae) from artificial structures and natural reefs in the Adriatic Sea. *Scientia Marina*, 74(4), 677-685. <https://doi.org/10.3989/scimar.2010.74n4677>
- Meiners-Mandujano, C., Fernández-Peralta, L., Faraj, A., & García-Cancela, R. (2018). Length-weight relationship of 15 deep-sea fish species (Actinopterygii) from the North-Western African continental slope. *Acta Ichthyologica et Piscatoria*, 48(2), 195-198. <https://doi.org/10.3750/AIEP/02042>
- Mendes, B., Fonseca, P., & Campos, A. (2004). Weight-length relationships for 46 fish species of the Portuguese west coast. *Journal of Applied Ichthyology*, 20(5), 355-361. <https://doi.org/10.1111/j.1439-0426.2004.00559.x>
- Miled-Fathali, N., Hamed, O., & Chakroun-Marzouk, N. (2019). Length-weight relationships of 22 commercial fish species from the Gulf of Tunis (Central Mediterranean Sea). *Cahiers de Biologie Marine*, 60, 541-546. <https://doi.org/10.21411/CBM.A.101E94A>
- Morey, G., Moranta, J., Massutí, E., Grau, A., Linde, M., Riera, F., & Morales-Nin, B. (2003). Weight-length relationships of littoral to lower slope fishes from the western Mediterranean. *Fisheries Research*, 62(1), 89-96. [https://doi.org/10.1016/S0165-7836\(02\)00250-3](https://doi.org/10.1016/S0165-7836(02)00250-3)
- Moutopoulos, D. K., & Stergiou, K. I. (2002). Length-weight and length-length relationships of fish species of the Aegean Sea (Greece). *Journal of Applied Ichthyology*, 18(3), 200-203. <https://doi.org/10.1046/j.1439-0426.2002.00281.x>
- Moutopoulos, D. K., Ramfos, A., Mouka, A., & Katselis, G. (2013). Length-weight relations of 34 fish species caught by small-scale fishery in Korinthiakos Gulf (Central Greece). *Acta Ichthyologica et Piscatoria*, 43(1), 57-64. <https://doi.org/10.3750/AIP2013.43.1.08>
- Muchlisin, Z. A., Musman, M., & Azizah, M. N. S. (2010). Length-weight relationships and condition factors of two threatened fishes, *Rasbora tawarensis* and *Poropuntius tawarensis*, endemic to Lake Laut Tawar, Aceh Province, Indonesia. *Journal of Applied Ichthyology*, 26(6), 949-953. <https://doi.org/10.1111/j.1439-0426.2010.01524.x>
- Ndiaye, W., Diouf, K., Samba, O., Ndiaye, P., & Panfili, J. (2015). The length-weight relationship and condition factor of white grouper (*Epinephelus aeneus*, Geoffroy Saint Hilaire, 1817) at the south-west coast of Senegal, West Africa. *International Journal of Advanced Research*, 3(3), 145-153.

- Özaydin, O., & Taskavak, E. (2006). Length-weight relationships for 47 fish species from Izmir Bay (eastern Aegean Sea, (Turkey). *Acta Adriatica*, 47(2), 211-216.
- Pauly, D. (1993). Fishbyte Section. Editorial. *Naga. ICLARM Quarterly*, 16, 26.
- Petrakis, G., & Stergiou, K. I. (1995). Weight-length relationships for 33 fish species in Greek waters. *Fisheries Research*, 21(3-4), 465-469. [https://doi.org/10.1016/0165-7836\(94\)00294-7](https://doi.org/10.1016/0165-7836(94)00294-7)
- Ricker, W. E. (1979). Growth rates and models. In W. S. Hoar, D. J. Randall, & J. R. Brett, (Eds.), *Fish Physiology, III, Bioenergetics and Growth* (pp 677-743). Academic Press.
- Ricker, W. E. (1968). *Methods for assessment of fish production in fresh waters*. IBP Handbook No. 3.
- Stergiou, K. I., & Moutopoulos, D. K. (2001). A review of length-weight relationships of fishes from Greek marine waters. *Naga. ICLARM Quarterly*, 24(1&2), 23-39.
- Tesch, F. W. (1971). Age and growth. In W. E. Ricker (Ed.), *Methods for assessment of fish production in fresh waters* (pp. 98-130). Blackwell Scientific Publications.
- Wootton, R. J. (2003). *Ecology of teleost fishes*. Chapman and Hall Ltd.