











## Compliance with Ethical Standards

### Authors' Contributions

The authors contributed equally to this paper.

### Conflict of Interest

The authors declare that there is no conflict of interest.

### Ethical Approval

For this type of study, formal consent was not required while the study conducted.

## REFERENCES

- Basin, M. (1995). Barcelona Resolution on the Environment and Sustainable Development in the Mediterranean Basin, Barcelona, June 1995.
- Bay-Nouailhat, A. (2007). Description of *Sciaena umbra*, available on line at <http://www.european-marine-life.org/34/sciaena-umbra.php>, consulted on 31 July 2019.
- Boglionne, C., Costa, C., Di Dato, P., Ferzini, G., Scardi, M., & Cataudella, S. (2003). Skeletal quality assessment of reared and wild sharpnose sea bream and pandora juveniles. *Aquaculture*, 227(1-4), 373-394. [https://doi.org/10.1016/S0044-8486\(03\)00515-5](https://doi.org/10.1016/S0044-8486(03)00515-5)
- Can, E. (2013). Effects of intensive and semi-intensive rearing on growth, survival, and V-shaped (Lordotic) skeletal deformities in juvenile gilthead sea bream (*Sparus aurata*). *Israeli Journal of Aquaculture-Bamidgeh*, 65, 1-7.
- Can, E., Karacalar, U., Saka, S., & Firat, K. (2012). Ozone disinfection of eggs from gilthead seabream *Sparus aurata*, sea bass *Dicentrarchus labrax*, red porgy, and common dentex *Dentex dentex*. *Journal of Aquatic Animal Health*, 24(2), 129-133. <https://doi.org/10.1080/08997659.2012.675925>
- Can, E., Saka, Ş., & Firat, K. (2010). Disinfection of gilthead sea bream (*Sparus aurata*), red porgy (*Pagrus pagrus*), and common dentex (*Dentex dentex*) eggs from Sparidae with different disinfectants. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*, 16(2), 299-306.
- Chao, L. (2015). *Sciaena umbra*. The IUCN Red List of Threatened Species 2015: e.T198707A83232286. <https://doi.org/10.2305/IUCN.UK.20154.RLTS.T198707A83232286.en>
- Chater, I., Romdhani, A., Ktari, M. H., & Mahe, K. (2012). Diet composition and feeding intensity of brown meagre *Sciaena umbra* Linnaeus, 1758 (Teleostei, Sciaenidae) in the Gulf of Tunis. *Proceedings of the 14th European Congress of Ichthyology*, Liège, Belgium. <https://archimer.ifremer.fr/doc/00117/22791/>
- Chatzifotis, S., Martin-Prat, A. V., Limberis, N., Papandroulakis, N., & Divanach, P. (2006). First data on growth of cultured brown meagre *Sciaena umbra* using diets with different protein and fat contents. *Fisheries Science*, 72(1), 83-88.
- Colla, S., Pranovi, F., Fiorin, R., Malavasi, S., & Picciulin, M. (2018). Using passive acoustics to assess habitat selection by the brown meagre *Sciaena umbra* in a northern Adriatic Sea mussel farm. *Journal of Fish Biology*, 92(5), 1627-1634. <https://doi.org/10.1111/jfb.13589>
- Divanach, P., & Kentouri, M. (2000). Hatchery techniques for specific diversification in Mediterranean finfish larviculture. *Cahiers Options Méditerranéennes*, 47, 75-87.
- Dou, S., Seikai, T., & Tsukamoto, K. (2000). Cannibalism in Japanese flounder juveniles, *Paralichthys olivaceus*, reared under controlled conditions. *Aquaculture*, 182(1-2), 149-159. [https://doi.org/10.1016/S0044-8486\(99\)00256-2](https://doi.org/10.1016/S0044-8486(99)00256-2)
- Engin, S., & Seyhan, K. (2009). Age, growth, sexual maturity and food composition of *Sciaena umbra* in the south-eastern Black Sea, Turkey. *Journal of Applied Ichthyology*, 25(1), 96-99. <https://doi.org/10.1111/j.1439-0426.2008.01173.x>
- Ergin, O., Tümer, S., & Yıldız, S. (2017). Chemical analysis of brown meager (*Sciaena umbra*) cephalides and traditional medicinal usage in urolithiasis. *Süleyman Demirel Üniversitesi Tıp Fakültesi Dergisi*, 24(1), 1-7.
- Hamzaçebi, S. (2020). A new species for aquaculture: White seabream (*Diplodus sargus*). *Journal of Anatolian Environmental and Animal Sciences*, 5(3), 390-396. <https://doi.org/10.35229/jaes.780711>
- Harmelin-Vivien, M., Cottalorda, J. M., Dominici, J. M., Harmelin, J. G., Le Diréach, L., & Ruitton, S. (2015). Effects of reserve protection level on the vulnerable fish species *Sciaena umbra* and implications for fishing management and policy. *Global Ecology and Conservation*, 3, 279-287. <https://doi.org/10.1016/j.gecco.2014.12.005>

- Kayim, M., Can, E., & Güner, Y. (2010). Is it possible to transform hatchery-reared abnormal juveniles of sea bass (*Dicentrarchus labrax* L. 1758) into normal individuals? *Asian Journal of Animal and Veterinary Advances*, 5(5), 327-338. <https://doi.org/10.3923/ajava.2010.327.338>
- Kolios, P., Kiritsis, S., & Katribusas, N. (1997). Larval-rearing and growout of the red porgy (*Pagrus pagrus*) in the Riopesca hatchery (Greece). *Hydrobiologia*, 358(1), 321-325. <https://doi.org/10.1023/A:1003132903785>
- Mediterranean to the Barcelona Convention (1995). Barcelona Convention for the Protection of the Marine Environment and the Coastal Region of the Mediterranean, 10 June 1995.
- Papadakis, I. E., Kentouri, M., Divanach, P., & Mylonas, C. C. (2013). Ontogeny of the digestive system of meagre *Argyrosomus regius* reared in a mesocosm, and quantitative changes of lipids in the liver from hatching to juvenile. *Aquaculture*, 388-391, 76-88. <https://doi.org/10.1016/j.aquaculture.2013.01.012>
- Papandroulakis, N., Divanach, P., Anastasiadis, P., & Kentouri, M. (2001). The pseudo-green water technique for intensive rearing of sea bream (*Sparus aurata*) larvae. *Aquaculture International*, 9(3), 205-216.
- Papandroulakis, N., Mylonas, C. C., Maingot, E., & Divanach, P. (2005). First results of greater amberjack (*Seriola dumerili*) larval rearing in mesocosm. *Aquaculture*, 250(1-2), 155-161. <https://doi.org/10.1016/j.aquaculture.2005.02.036>
- Roo, F. J., Hernández-Cruz, C. M., Socorro, J. A., Fernández-Palacios, H., & Izquierdo, M. S. (2010). Advances in rearing techniques of *Pagrus pagrus* (Linnaeus, 1758): Comparison between intensive and semi-intensive larval rearing systems. *Aquaculture Research*, 41(3), 433-449. <https://doi.org/10.1111/j.1365-2109.2009.02244.x>
- Russo, T., Scardi, M., Boglione, C., & Cataudella, S. (2010). Application of the Self-Organizing Map to the study of skeletal anomalies in aquaculture: The case of dusky grouper (*Epinephelus marginatus* Lowe, 1834) juveniles reared under different rearing conditions. *Aquaculture*, 315(1-2), 69-77. <https://doi.org/10.1016/j.aquaculture.2010.11.030>
- Saka, Ş., Firat, K., & Kamaci, H. O. (2001). The development of European sea bass (*Dicentrarchus labrax* L., 1758) eggs in relation to temperature. *Turkish Journal of Veterinary and Animal Sciences*, 25(2), 139-147.
- Shields, R. J., Gara, B., & Gillespie, M. J. (1999). A UK perspective on intensive hatchery rearing methods for Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture*, 176(1-2), 15-25. [https://doi.org/10.1016/S0044-8486\(99\)00046-0](https://doi.org/10.1016/S0044-8486(99)00046-0)
- Valles, R., & Estevez, A. (2013). Light conditions for larval rearing of meagre (*Argyrosomus regius*). *Aquaculture*, 376, 15-19. <https://doi.org/10.1016/j.aquaculture.2012.11.011>
- Zaiss, M. M., Papadakis, I. E., Maingot, E., Divanach, P., & Mylonas, C. C. (2006). Ontogeny of the digestive tract in shi drum (*Umbrina cirrosa* L.) reared using the mesocosm larval rearing system. *Aquaculture*, 260(1-4), 357-368. <https://doi.org/10.1016/j.aquaculture.2006.06.001>