

- Santini, F., Carnevale, G., & Sorenson, L. (2014). First multi-locus timetree of seabreams and porgies (Percomorpha: Sparidae). *Italian Journal of Zoology*, 81, 55-71. <https://doi.org/10.1080/11250003.2013.878960>
- Sarı, E., & Çağatay, M. N. (2001). Distributions of heavy metals in the surface sediments of the Gulf of Saros, NE Aegean Sea. *Environment International*, 26, 169-173. [https://doi.org/10.1016/s0160-4120\(00\)00097-0](https://doi.org/10.1016/s0160-4120(00)00097-0)
- Saygılı, B., İşmen, A., & İhsanoğlu, M. A. (2016a). Age and growth of blotched picarel (*Spicara maena* Linnaeus, 1758) in the Sea of Marmara and Northern Aegean Sea. *Ege Journal of Fisheries & Aquatic Sciences*, 33(2), 143-149. <https://doi.org/10.12714/egejfas.2016.33.2.08>
- Saygılı, B., İşmen, A., & İhsanoğlu, M. A. (2016b). Differences in the otolith shape between the Marmara and Northern Aegean Sea stocks of blotched picarel (*Spicara maena* Linnaeus, 1758). *Journal of the Black Sea/Mediterranean Environment*, 22, 137-148.
- Soykan, O., İlkyaz, A. T., Metin, G., & Kınacıgil, H. T. (2010). Growth and reproduction of blotched picarel (*Spicara maena* Linnaeus, 1758) in the Central Aegean Sea Turkey. *Turkish Journal of Zoology*, 34, 453-459. <https://doi.org/10.3906/zoo-0903-29>
- Turley, C. M., Bianchi, M., Christaki, U., Conan, P., Harris, J. R. W., Psarra, S., Ruddy, G., Stutt, E. D., Tselepides, A., & Van Wambeke, F. (2000). The relationship between primary producers and bacteria in an oligotrophic sea - The Mediterranean and biogeochemical implications. *Marine Ecology Progress Series*, 193, 11-18. <https://doi.org/10.3354/meps193011>
- Wootton, R. J. (1990). *Ecology of teleost fish*. Chapman and Hall.