





















- Trushenski, J., Schwarz, M., Lewis, H., Laporte, J., Delbos, B., Takeuchi, R., & Sampaio, L. A. (2011). Effect of replacing dietary fish oil with soybean oil on production performance and fillet lipid and fatty acid composition of juvenile cobia *Rachycentron canadum*. *Aquaculture Nutrition*, 17(2), 437-447. <https://doi.org/10.1111/j.1365-2095.2010.00779.x>
- Webster, C. D., Rawles, S. D., Koch, J. F., Thompson, K. R., Kobayashi, Y., Gannam, A. L., Twibel, R. G., & Hyde, N. M. (2016). Bio-Ag reutilization of distiller's dried grains with solubles (DDGS) as a substrate for black soldier fly larvae, *Hermetia illucens*, along with poultry by-product meal and soybean meal, as total replacement of fish meal in diets for Nile tilapia, *Oreochromis niloticus*. *Aquaculture Nutrition*, 22(5), 976-988. <https://doi.org/10.1111/anu.12316>
- Yıldız, M., & Şener, E. (2004). Karadeniz Bölgesi'ndeki gökkuşuğu alabalığı (*Oncorhynchus mykiss*) ve deniz levreği (*Dicentrarchus labrax*) yetiştiriciliği yapan işletmelerin yapısal analizi ve biyo-teknolojik özellikleri [Structural and biotechnological properties of rainbow trout (*Oncorhynchus mykiss*) and seabass (*Dicentrarchus labrax*) farms in the Black Sea region]. *İstanbul Üniversitesi Veteriner Fakültesi Dergisi*, 29(2), 241-252.
- Yılmaz, S., & Ergün, S. (2013). Chickweed (*Stellaria media*) leaf meal as a feed ingredient for tilapia (*Oreochromis mossambicus*). *Journal of Applied Aquaculture*, 25(4), 329-336. <https://doi.org/10.1080/10454438.2013.851531>