



















- Aktar, S., & Cebe G. E. (2010). General specifications, using areas of algae and their importance on pharmacy. *Ankara Üniversitesi Eczacılık Fakültesi Dergisi*, 39(3), 237-264. <https://doi.org/10.1501/Eczfak.0000000568>
- Alçay, A. Ü., Bostan, K., Dinçel, E., & Varlık, C. (2017). *Alglerin insan gıdası olarak kullanımı* [Algae as a food source for humans]. *Aydın Gastronomy*, 1(1), 47-59.
- Bligh, E. G., & Dyer, W. J. (1959). A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology*, 37(8), 911-917. <https://doi.org/10.1139/o59-099>
- Caf, F., Yılmaz, Ö., Durucan F., & Özdemir, N. Ş. (2015). Biochemical components of three marine macroalgae (*Padina pavonica*, *Ulva lactuca* and *Taonia atomaria*) from the Levantine Sea coast of Antalya, Turkey. *Journal of Biodiversity and Environmental Sciences*, 6(4), 401-411.
- Carvalho, J., Ribeiro, A., Castro, J. D. F. de, Vilarinho, C., & Castro, F. (2011). Biodiesel production by microalgae and macroalgae from north littoral Portuguese coast. *Proceedings of the 1st International Conference on Wastes: Solutions, Treatments and Opportunities*, Portugal. <https://core.ac.uk/download/pdf/55615137.pdf>
- Colombo, M. L., Rise, P., & Giavarini, F. (2006). Marine microalgae as sources of polyunsaturated fatty acids. *Plant Foods for Human Nutrition*, 61, 64-69. <https://doi.org/10.1007/s1130-006-0015-7>
- Cowing, B. E., & Saker, K. E. (2001). Polyunsaturated fatty acids and epidermal growth factor receptor/mitogen-activated protein kinase signaling in mammary cancer. *The Journal of Nutrition*, 131(4), 1125-1128. <https://doi.org/10.1093/jn/131.4.1125>
- El-Sheekh, M. M., Osman, M. E. H., Dyab, M. A., Amer, M. S. (2006). Production and characterization of antimicrobial active substance from the cyanobacterium *Nostoc muscorum*. *Environmental Toxicology and Pharmacology*, 21(1), 42-50. <https://doi.org/10.1016/j.etap.2005.06.006>
- EL-Wakf, A. M., Ebraheem, H. A., Serag, H. A., Hassan, H. A., & Gumaih, H. S. (2010). Association between inflammation and the risk of cardiovascular disorders in atherogenic male rats: Role of virgin and refined olive oil. *Journal of American Science*, 6(12), 807-817.
- Ganesan, K., Suresh Kumar, K., & Subba Rao, P. V. (2011). Comparative assessment of antioxidant activity in three edible species of green seaweed, *Enteromorpha* from Okha, Northwest coast of India. *Innovative Food Science & Emerging Technologies*, 12(1), 73-78. <https://doi.org/10.1016/j.ifset.2010.11.005>
- Ganesan, K., Suresh Kumar, K., Subba Rao, P. V., Tsukui, Y., Bhaskar, N., Hosokawa, M., & Miyashita, K. (2014). Studies on chemical composition of three species of *Enteromorpha*. *Biomedicine & Preventive Nutrition*, 4(3), 365-369. <https://doi.org/10.1016/j.bionut.2014.04.001>
- Gümüş, G. (2006). *Deniz marulunun kimyasal kompozisyonunun araştırılması*. [Yüksek Lisans Tezi. Ege Üniversitesi].
- Guner, H., & Aysel, V. (1989). *Tohumuz bitkiler I. cilt: Algler*. Ege Üniversitesi Basımevi.
- Guschina, I. A., & Harwood, J. L. (2006). Lipids and lipid metabolism in eukaryotic algae. *Progress in Lipid Research*, 45(2), 160-186. <https://doi.org/10.1016/j.plipres.2006.01.001>
- Harwood, J. L., & Guschina, I. A. (2009). The versatility of algae and their lipid metabolism. *Biochimie*, 91(6), 679-684. <https://doi.org/10.1016/j.biochi.2008.11.004>
- HMSO. (1994). *Nutritional aspects of cardiovascular disease*. Report on health and social subjects no. 46. London, UK.
- Holdt, S. L., & Kraan S. (2011). Bioactive compounds in seaweed: functional food applications and legislation. *Journal of Applied Phycology*, 23, 543-597. <https://doi.org/10.1007/s10811-010-9632-5>
- Ichibara, K., Shibahara, A., Yamamoto, K., & Nakayama, T. (1996). An improved method for rapid analysis of the fatty acids of glycerolipids. *Lipids*, 31(5), 535-539. <https://doi.org/10.1007/bf02522648>
- Kandasamy, G., Karuppiah, S. K., & Subba Rao, P. V. (2012). Salt- and pH-induced functional changes in protein concentrate of edible green seaweed *Enteromorpha* species. *Fisheries Science*, 78, 169-176. <https://doi.org/10.1007/s12562-011-0423-y>
- Lewis, M. A., & Wang, W. (1997). Water quality and aquatic plants. In W. Wang, W. R. Lower, J. W. Gorsuch, & J. S. Hughes (Eds.), *Plants for environmental studies* (pp. 141-175). CRC Lewis Publishers.
- Mchugh, D. J. (2003). *A guide to the seaweed industry*. FAO Fisheries Technical Paper, No. 441. Rome. 105p.
- Spolaore, P., Joannis-Cassan, C., Duran, E., & Isambert, A. (2006). Commercial applications of microalgae. *Journal of Bioscience and Bioengineering*, 101(2), 87-96. <https://doi.org/10.1263/jbb.101.87>
- Tabudravu, J. N., Gangaiya, P., Sotheeswaran, S., & South, G. R. (2002). *Enteromorpha flexuosa* (Wulfen) J. Agardh (Chlorophyta: Ulvales)--evaluation as an indicator of heavy metal contamination in a tropical estuary. *Environmental Monitoring and Assessment*, 75(2), 201-213. <https://doi.org/10.1023/A:1014439931466>
- Ulbricht, T. L. V., & Southgate, D. A. T. (1991). Coronary heart disease: seven dietary factors. *The Lancet*, 338, 985-992. [https://doi.org/10.1016/0140-6736\(91\)91846-m](https://doi.org/10.1016/0140-6736(91)91846-m)

- Wahbeh, M. I. (1997). Amino acid and fatty acid profiles of four species of macroalgae from Aqaba and their suitability for use in fish diets. *Aquaculture*, 159(1-2), 101-109. [https://doi.org/10.1016/S0044-8486\(97\)00183-X](https://doi.org/10.1016/S0044-8486(97)00183-X)
- Weihrach, J. L., Posati, L. P., Anderson, B. A., & Exler, J. (1975). Lipid conversion factors for calculating fatty acid contents of foods. *Journal of the American Oil Chemists' Society*, 54(1), 36-40. <https://doi.org/10.1007/BF02671370>
- Zeybek, N., Zeybek, U., & Sayginer, B. (2003). *Farmasötik botanik*. Meta Basimevi.