

Compliance With Ethical Standards

Conflict of Interest

The author declares that there is no conflict of interest.

Ethical Approval

For this type of study, formal consent is not required.

REFERENCES

- Abbasi, K., Mouludi-Saleh, A., Eagderi, S., & Sarpanah, A. (2019). Length-weight relationship and condition factor of eight species of the genera *Capoeta*, *Garra*, *Chondrostoma*, *Schizothorax* and *Paraschistura* from Iranian inland waters. *Iranian Journal of Ichthyology*, 6(4), 26-270. <https://doi.org/10.22034/iji.v6i4.432>
- Altın, A., Ayyıldız, H., Kale, S., & Alver, C. (2015). Length-weight relationships of forty-nine fish species from shallow waters of Gökçeada Island, northern Aegean Sea. *Turkish Journal of Zoology*, 39(5), 971-975. <https://doi.org/10.3906/zoo-1412-15>
- Aydın, M. (2020). First report of *Symphodus melops* (Linnaeus, 1758) with maximum length in the Black Sea. *Marine Science and Technology Bulletin*, 9(2), 125-129. <https://doi.org/10.33714/masteb.741985>
- Bagenal, T. B., & Tesch, F. W. (1978). Age and growth. In T. Bagenal (Ed.), *Methods for assessment of fish production in fresh waters* (pp. 101-136). 3rd ed. IBP Handbook No. 3, Blackwell Science Publications.
- Bilecenoğlu, M., Kaya, M., Cihangir, B., & Çiçek, E. (2014). An updated checklist of the marine fishes of Turkey. *Turkish Journal of Zoology*, 38(6), 901-929. <https://doi.org/10.3906/zoo-1405-60>
- Bilge, G., Yapıcı, S., Filiz, H., & Cerim, H. (2014). Weight-length relations for 103 fish species from the southern Aegean Sea, Turkey. *Acta Ichthyologica et Piscatoria*, 44(3), 263-269. <https://doi.org/10.3750/AIP2014.44.3.11>
- Cengiz, Ö., İşmen, A., Özekinci, U., & Öztekin, A. (2011). An investigation on fish fauna of Saros Bay (Northern Aegean Sea). *Afyon Kocatepe University Journal of Sciences*, 11(1), 31-37.
- Cengiz, Ö., Öztekin, A., & Özekinci, U. (2012). An investigation on fishes spreading along the coasts of Gallipoli Peninsula and Dardanelles (North-eastern Mediterranean, Turkey). *Firat University Journal of Science*, 24, 47-55.
- Cengiz, Ö. (2020). A study on maximum length record of saddled seabream (*Oblada melanura* Linnaeus, 1758) caught off Gökçeada Island (Northern Aegean Sea, Turkey). *Marine Science and Technology Bulletin*, 9(1), 58-61. <https://doi.org/10.33714/masteb.691478>
- Cengiz, Ö., & Paruğ, Ş. Ş. (2020). A new record of the rarely reported grey triggerfish (*Balistes capriscus* Gmelin, 1789) from Northern Aegean Sea (Turkey). *Marine and Life Sciences*, 2(1), 1-4.
- Cengiz, Ö. (2021). Opercular girth, maximum girth and total length relationships for eight fish species from the Saros Bay (northern Aegean Sea, Turkey). *The Palawan Scientist*, 13(2), *In press*.
- Cengiz, Ö., & Paruğ, Ş. Ş. (2021). Growth parameters of blotched picarel (*Spicara maena* Linnaeus, 1758) from Saros Bay (Northern Aegean Sea, Turkey). *Acta Natura et Scientia*, 2(1), 40-48. <https://doi.org/10.29329/actanatsci.2021.314.7>
- Deekae, S. N., & Abowei, J. F. N. (2010). *Macrobrachium macrobrachion* (Herklots, 1851) length-weight relationship and Fulton's condition factor in Luubara creek, Ogoni land, Niger Delta, Nigeria. *International Journal of Animal and Veterinary Advances*, 2(4), 155-162.
- Dimitriadis, C., & Fournari-Konstantinidou, I. (2018). Length weight relations for 20 fish species (Actinopterygii) from the Southern Ionian Sea, Greece. *Acta Ichthyologica et Piscatoria*, 48(4), 415-417. <https://doi.org/10.3750/AIEP/02466>
- Dutta, S., Maity, A., Chanda, A., Akhand, A., & Hazra, S. (2012). Length weight relationship of four commercially important marine fishes of Northern Bay of Bengal, West Bengal, India. *Journal of Applied Environmental Biological Sciences*, 2, 52-58.
- Eagderi, S., Mouludi-Saleh, A., & Cicek, E. (2020). Length-weight relationship of ten species of Leuciscinae sub-family (Cyprinidae) from Iranian inland waters. *International Aquatic Research*, 12(2), 133-136. [https://doi.org/10.22034/iar\(20\).2020.1891648.1004](https://doi.org/10.22034/iar(20).2020.1891648.1004)
- Froese, R. (2006). Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22(4), 241-253. <https://doi.org/10.1111/j.1439-0426.2006.00805.x>
- Froese, R., & Pauly, D. (Eds) (2020). FishBase. World Wide Web electronic publication. www.fishbase.org, version (12/2020). [Accessed 09 March 2021]
- Gürkan, Ş., Bayhan, B., Akcınar, S. C., & Taskavak, E. (2010). Length-weight relationship of fish from shallow waters of Candarli Bay (North Aegean Sea, Turkey). *Pakistan Journal of Zoology*, 42(4), 495-498.
- İlhan, D. U., Akalın, S., Tosunoğlu, Z., & Özyayın, O. (2008). Length-weight relationships of five *Symphodus* species (Pisces: Perciformes) from İzmir Bay, Aegean Sea. *Ege University Journal of Fisheries and Aquatic Sciences*, 25(3), 245-246.

- Kalaycı, F., Samsun, N., Samsun, O., Şahin, C., & Dalgıç, G. (2007). The determination of biologic characteristics and meat yield of *Symphodus tinca*, Linnaeus, 1758 (Labridae) caught as discard from Sinop area. *Proceedings of the XIV. National Fisheries Symposium*, Turkey, pp. 154-155.
- Kasapoglu, N., Cankiriligil, E. C., Firidin, S., & Cakmak, E. (2016). A preliminary study on some biological characters of East Atlantic peacock wrasse, *Symphodus tinca*, in the Black Sea, Turkey. *Journal of the Black Sea/Mediterranean Environment*, 22(3), 289-294.
- Karakulak, F. S., Erk, H., & Bilgin, B. (2006). Length-weight relationships for 47 coastal fish species from the northern Aegean Sea, Turkey. *Journal of Applied Ichthyology*, 22(4), 274-278. <https://doi.org/10.1111/j.1439-0426.2006.00736.x>
- Keskin, Ç., & Gaygusuz, Ö. (2010). Length-weight relationships of fishes in shallow waters of Erdek Bay (Sea of Marmara, Turkey). *European Journal of Biology*, 69(2), 87-94.
- Keskin, Ç., & Ünsal, N. (1998). The fish fauna of Gökçeada Island, NE Aegean Sea, Turkey. *Italian Journal of Zoology*, 65(Suppl.), 299-302.
- Le Cren, E. D. (1951). The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology*, 20(2), 201-219. <https://doi.org/10.2307/1540>
- Maravelias, C. D., & Papaconstantinou, C. (2006). Geographic, seasonal and bathymetric distribution of demersal fish species in the eastern Mediterranean. *Journal of Applied Ichthyology*, 22(1), 35-42. <https://doi.org/10.1111/j.1439-0426.2006.00695.x>
- Mater, S., Kaya, M., & Bilecenoğlu, M. (2003). *Türkiye deniz balıkları atlası* (3. Ed.). Ege Üniversitesi Su Ürünleri Fakültesi Yayınları, No. 68.
- Metaxas, D. A. (1973). Air-sea interaction in the Greek seas and resultant Etesian characteristics. Tech Rep 5, University of Ioannina, pp. 1-32.
- Miled-Fathalli, N., Hamed, O., & Chakroun-Marzouk, N. (2019). Length weight relationships of 22 commercial fish species from the Gulf of Tunis (Central Mediterranean Sea). *Cahiers de Biologie Marine*, 60(6), 541-546. <https://doi.org/10.21411/CBM.A.101E94A>
- Muchlisin, Z. A., Musman, M., & Azizah, M. N. S. (2010). Length-weight relationships and condition factors of two threatened fishes, *Rasbora tawarensis* and *Poropuntius tawarensis*, endemic to Lake Laut Tawar, Aceh Province, Indonesia. *Journal of Applied Ichthyology*, 26(6), 949-953. <https://doi.org/10.1111/j.1439-0426.2010.01524.x>
- Nival, S. (1966). Contribution a l'étude biologique de *Symphodus rostratus* Bl. (Labridae) en rade de Villefranche-sur-mer. *Vie Milieu*, 17(3A), 1273-1315.
- Ndiaye, W., Diouf, K., Samba, O., Ndiaye, P., & Panfili, J. (2015). The length-weight relationship and condition factor of white grouper (*Epinephelus aeneus*, Geoffroy Saint Hilaire, 1817) at the south-west coast of Senegal, West Africa. *International Journal of Advanced Research*, 3(3), 145-153.
- Olson, D. B., Kourafalou, V. H., Johns, W. E., Samuels, G., Veneziani, M. (2007). Aegean surface circulation from a satellite-tracked drifter array. *Journal of Physical Oceanography*, 37(7), 1898-1917. <https://doi.org/10.1175/JPO3028.1>
- Onay, H. (2021). Length-weight relationships of four *Symphodus* species (Actinopterygii: Perciformes: Labridae) from Eastern Black Sea (Turkey). *Marine Science and Technology Bulletin*, 10(3), 228-233. <https://doi.org/10.33714/masteb.824561>
- Özaydın, O., Uçkun, D., Akalın, S., Leblebici, S., & Tosunoğlu, Z. (2007). Length-weight relationships of fishes captured from Izmir Bay, Central Aegean Sea. *Journal of Applied Ichthyology*, 23(6), 695-696. <https://doi.org/10.1111/j.1439-0426.2007.00853.x>
- Pallaoro, A., & Jardas, I. (2003). Some biological parameters of the peacock wrasse, *Symphodus (Crenilabrus) tinca* (L. 1758) (Pisces: Labridae) from the middle Eastern Adriatic (Croatian coast). *Scientia Marina*, 67(1), 33-41. <https://doi.org/10.3989/scimar.2003.67n133>
- Parenti, P., & Randall, J. E. (2011). Checklist of the species of the families Labridae and Scaridae: An update. *Smithiana Bulletin*, 13, 29-44.
- Paruğ, Ş., & Cengiz, Ö. (2020). The maximum length record of the white seabream (*Diplodus sargus* Linnaeus, 1758) for the Aegean Sea. *Acta Natura et Scientia*, 1(1), 96-108. <https://doi.org/10.29329/actanatsci.2020.313.11>
- Petrakis, G., & Stergiou, K. I. (1995). Weight-length relationships for 33 fish species in Greek waters. *Fisheries Research*, 21(3-4), 465-469. [https://doi.org/10.1016/0165-7836\(94\)00294-7](https://doi.org/10.1016/0165-7836(94)00294-7)
- Raposeiro, P. M. V. M., & Azevedo, J. M. N. (2009). Reproductive biology of *Symphodus mediterraneus* (Teleostei, Labridae) in the Azores. *Marine Ecology*, 30(Suppl.1), 175-180. <https://doi.org/10.1111/j.1439-0485.2009.00309.x>
- Ricker, W. E. (1968). *Methods for assessment of fish production in fresh waters*. IBP Handbook No. 3.
- Škeljo, F., Brčić, J., Vuletin, V., & Ferri, F. (2015). Age and growth of the axillary wrasse, *Symphodus mediterraneus* (L.) from the eastern Adriatic Sea. *Marine Biology Research*, 11(7), 780-784. <https://doi.org/10.1080/17451000.2015.1016963>

- Stergiou, K. I., & Moutopoulos, D. K. (2001). A review of length-weight relationships of fishes from Greek marine waters. *Naga, The ICLARM Quarterly*, 24(1&2), 23-39.
- Tesch, F. W. (1971). Age and growth. In W. E. Ricker (Ed.), *Methods for assessment of fish production in fresh waters* (pp. 98-130). Blackwell Scientific Publications.
- Theocharis, A., Balopoulos, E., Kioroglou, S., Kontoyiannis, H., & Iona, A. (1999). A synthesis of the circulation and hydrography of the South Aegean Sea and the Straits of the Cretan Arc (March 1994-January 1995). *Progress In Oceanography*, 44(4), 469-509. [https://doi.org/10.1016/S0079-6611\(99\)00041-5](https://doi.org/10.1016/S0079-6611(99)00041-5)
- Torres, M. A., Ramos, F., & Sobrino, I. (2012). Length-weight relationships of 76 fish species from the Gulf of Cadiz (SW Spain). *Fisheries Research*, 127-128, 171-175. <https://doi.org/10.1016/j.fishres.2012.02.001>
- Valle, C., Bayle, J. T., & Ramos, A. A. (2003). Weight-length relationships for selected fish species of the western Mediterranean Sea. *Journal of Applied Ichthyology*, 19(4), 261-262. <https://doi.org/10.1046/j.1439-0426.2003.00492.x>
- Whitehead, P. J. P., Bauchot, M. L., Hureau, J. C., Nielsen, J., & Tortonese, E. (Eds.) (1986). *Fishes of the north-eastern Atlantic and the Mediterranean*. UNESCO.
- Wootton, R. J. (2003). *Ecology of teleost fishes*. Chapman and Hall Ltd.
- Yıldırım, A., Erdoğan, O., Türkmen, M., & Demir, B. C. (1999). The investigation of some reproduction characteristics of the *Alburnoides bipunctatus* fasciitis (Nordman, 1840) living in Oltu Stream of Coruh River (Artvin, Turkey). *Turkish Journal of Veterinary and Animal Sciences*, 23(Ek Sayı 4), 163-168.
- Zodiatis, G., & Balopoulos, E. (1993). Structure and characteristics of fronts in the North Aegean Sea. *Bollettino di Oceanologia Teorica ed Applicata*, 11(2), 113-124.