










Impact of Different Nutrient Enrichment Concentrations on the Growth of Microalga *Nannochloropsis* sp. (Monodopsidaceae) Culture

Norsheen B. Sanuddin¹  • Melodina D. Hairol¹  • Cherry T. Nian¹  • Rizal Jhunn F. Robles¹  •
Hadjiran A. Illud¹  • Jubail S. Muyong¹  • Jamrun H. Ebbah¹  • Jurmin H. Sarri¹  

¹ Mindanao State University-Tawi-Tawi College of Technology and Oceanography, College of Fisheries, Sanga-Sanga, Bongao, 7500 Tawi-Tawi, Philippines, nsanuddin32@gmail.com, melodinahairol@gmail.com, cherrynian2022@gmail.com, rizaljhunnr@gmail.com, illudhadjiran@gmail.com, muyongjubail2@gmail.com, ebbahjamrun@gmail.com, jurslink@gmail.com

✉ Corresponding Author: jurslink@gmail.com

Please cite this paper as follows:

Sanuddin, N. B., Hairol, M. D., Nian, C. T., Robles, R. J. F., Illud, H. A., Muyong, J. S., Ebbah, J. H., & Sarri, J. H. (2023). Impact of Different Nutrient Enrichment Concentrations on the Growth of Microalga *Nannochloropsis* sp. (Monodopsidaceae) Culture. *Acta Natura et Scientia*, 4(1), 87-93. <https://doi.org/10.29329/actanatsci.2023.353.09>

ARTICLE INFO

Article History

Received: 02.05.2023

Revised: 10.06.2023

Accepted: 20.06.2023

Available online: 22.06.2023

Keywords:

Cell density

Growth

Microalgae

Nannochloropsis sp.

Nutrient enrichment

A B S T R A C T

Microalgae consist of unicellular algal species that can produce and accumulate a wide variety of biomolecules. In order to maintain a high cell density in a continuous phototrophic culture of algae, the nutrient can serve as the most important factor in enhancing cell density. In this study, the effect of different concentrations of nutrients on the cell density of microalga *Nannochloropsis* sp. cultured in the mega plastic box (with 50 L capacity) was investigated. Four groups of treatment with four replicates were tested: group A (including 5 g L⁻¹ of ferric chloride, ammonium phosphate, and Urea), group B (including 10 g L⁻¹ of ferric chloride, ammonium phosphate, and urea), group C (including 15 g L⁻¹ of ferric chloride, ammonium phosphate, and urea), and group D (including 15 g L⁻¹ of ferric chloride, ammonium phosphate, and urea with 40 g L⁻¹ of cow manure). Results revealed that group C and group D achieved maximum density on day three as 86.39×10⁶ cell mL⁻¹ and 85.59×10⁶ cell mL⁻¹, respectively, which were significantly (p≤0.05) higher than the cell density of groups A (58.01×10⁶ cell mL⁻¹) and group B (70.67×10⁶ cell mL⁻¹). Additionally, the increasing specific growth rate (SGR) of *Nannochloropsis* sp. cultured was obtained in group D at 0.308 day⁻¹ after the culture period. From the result of the study, it is concluded that the concentrations of 15 g L⁻¹ ferric chloride, ammonium phosphate, and urea (group C) and 15 g L⁻¹ ferric chloride, ammonium phosphate, and urea combined with 40 g L⁻¹ cow manure (group D) are capable of increasing cell density growth of microalga *Nannochloropsis* sp. cultured in a mega plastic box.

INTRODUCTION

Microalgae can be used in many biotechnological applications, including health foods, feeds, colorants, bioenergy, and pharmaceuticals (Pulz & Gross, 2004; Mortensen, 2006; Chu, 2012; Dixit & Suseela, 2013). Microalgal compounds are particularly valuable for finding new medicines that treat viral infections, cancers, and bacteria and fungi showing resistance to antibiotic treatments (Skulberg, 2000; Huleihel et al., 2001). In addition, various microalgae contain DHA (docosahexaenoic acid), which contributes to heart health in adults and brain development in babies (Pulz & Gross, 2004). In addition to their role in photosynthesis, microalgal pigments have a wide range of biological properties, including antioxidants (Hejazi et al., 2004; Chidambara-Murthy et al., 2005), anti-obesity, anti-angiogenic, anti-cancer, and neuroprotective properties (Ciccone et al., 2013), as well as an anti-inflammatory (García-González & Ochoa, 1999; Guzman et al., 2001). Growing concern about human nutrition has led to controversy regarding microalgae in poultry and fish (Souza et al., 2021). Furthermore, researchers examined how microalgae can be cultivated rapidly and produced in large quantities (Duong et al., 2012; Leu & Boussiba, 2014; Mohan et al., 2015). Various culturing systems have been used to cultivate microalgae at large scales under controlled conditions, including polyethylene sleeves, outdoor ponds, and tubular bioreactors (Lebeau & Robert, 2003; Benner et al., 2022). Moreover, microalgal growth is influenced by environmental parameters such as temperature, light intensity, and photoperiod in culture systems (George et al., 2014). There are a number of factors that affect microalgae growth and biochemical composition, including nutrients (Juneja et al., 2013; Barkia et al., 2019). The manure of terrestrial animals can have significant effects on plants, including macroalgae (Jjemba, 2002; Bogaard et al., 2007; Giwa, 2017). However, the studies related to the production of microalgae with manure and other inorganic nutrients are very limited. Thus, this study examines the effect of different concentrations of nutrients along with cow manure on the growth of *Nannochloropsis* sp. culture.

MATERIAL AND METHODS

Study Area

The experiment was conducted at the Multi-species Hatchery, situated in the College of Fisheries, Mindanao State University-Tawi-Tawi College of Technology and Oceanography (MSU-TCTO), Sanga-Sanga, Bongao, Tawi-Tawi, Philippines (05°02'3.8" N 119°44'5.0" E) on March 23-27, 2022.

Culture Condition

Mega plastic boxes with 50 L volume were used in *Nannochloropsis* sp. experiment. Chlorinated seawater was used in this study and filtered through a filter bag before being transported to 16 mega boxes containing 45 liters each. After the 16 mega boxes were filled with chlorinated seawater, starter green microalgae of *Nannochloropsis* sp. with 5 L volume inoculated into the culture container. Mega plastic boxes were then mixed with four different nutrient concentrations randomly (Table 1) with four replicates for each treatment. The mega plastic boxes were covered with transparent acetate film to avoid contamination, and fluorescent lamps (Ecolum LED) were used as artificial lighting for the cultures (Figure 1).

Analysis

A daily sample of *Nannochloropsis* sp. culture was taken for cell counting and analysis. The number of cells was counted by using a hemocytometer, and contamination was investigated under a microscope. Specific growth rates (μ) were calculated by following the formula (Durmaz & Erbil, 2017);

$$\mu = \frac{\ln(N_2) - \ln(N_1)}{t_2 - t_1} \quad (1)$$

where: N_2 is the biomass cell number at the time (t_2). N_1 is the beginning biomass cell number at the time (t_1).

IBM SPSS software version 20 was used to analyze the significance of differences between the growth using one-way analysis of variance (ANOVA). Post hoc (Duncan) was used to rank the means.

Table 1. Different concentrations of nutrients of ferric chloride, ammonium phosphate, urea, and cow manure that used in four different experimental treatment groups

Treatment	Ferric Chloride (g L ⁻¹)	Ammonium Phosphate (g L ⁻¹)	Urea (g L ⁻¹)	Cow Manure (g L ⁻¹)
Group A	5	5	5	0
Group B	10	10	10	0
Group C	15	15	15	0
Group D	15	15	15	40




Figure 1. Experimental set-up

RESULTS

The initial cell density of *Nannochloropsis* sp. was arranged as 3.20×10^6 cell mL⁻¹ (Figure 2). The microalga *Nannochloropsis* sp. was enriched with different concentrations of nutrients. All of the experimental groups reached the maximum cell density on the 3rd day of culture. The maximum cell density of groups C and D on 3rd day of the culture period was 86.39×10^6 cell mL⁻¹ and 85.59×10^6 cell mL⁻¹, respectively, which was significantly ($p \leq 0.05$) higher than the cell density of groups A and B at 58.01×10^6 cell mL⁻¹ and 70.67×10^6 cell mL⁻¹, respectively. The lag phase was observed on the first day for all the group treatments. After the first day, the cell density of *Nannochloropsis* sp. in group A increased rapidly from 29.55×10^6 cell mL⁻¹, reaching 50.01×10^6 cell mL⁻¹, while in group B, the cell density increased from 27.59×10^6

cell mL⁻¹ to 70.67×10^6 cell mL⁻¹ by day 3 without any apparent lag phase. Moreover, the cell density in group C was increased from 27.97×10^6 cell mL⁻¹ to 86.39×10^6 cell mL⁻¹, while in group D it was increased from 31.05×10^6 cell mL⁻¹ to 85.59×10^6 cell mL⁻¹ at the 3rd day. Furthermore, the specific growth rate (SGR) of *Nannochloropsis* sp. culture is shown in Figure 3. The SGR of *Nannochloropsis* sp. culture in group D (0.308 ± 0.003 day⁻¹) was increased than the SGR in group A (0.170 ± 0.011 day⁻¹), B (0.173 ± 0.011 day⁻¹), and C (0.188 ± 0.019 day⁻¹) at the end of culture period.

DISCUSSION

The primary aim of producing phototrophic organisms is to maintain a high cell density in a continuous culture. In the present study, green microalga *Nannochloropsis* sp. was investigated in




Figure 2. Cell densities of different concentrations of nutrient enrichment. A = (5 g L⁻¹ of ferric chloride, ammonium phosphate, and Urea); B = (10 g L⁻¹ of ferric chloride, ammonium phosphate, and Urea); C = (15 g L⁻¹ of ferric chloride, ammonium phosphate, and Urea); D (15 g L⁻¹ of ferric chloride, ammonium phosphate, and Urea with 40 g L⁻¹ of cow manure), N=16.




Figure 3. Specific growth rate of different concentrations of nutrient enrichment. A = (5 g L⁻¹ of ferric chloride, ammonium phosphate, and Urea); B = (10 g L⁻¹ of ferric chloride, ammonium phosphate, and Urea); C = (15 g L⁻¹ of ferric chloride, ammonium phosphate, and Urea); D (15 g L⁻¹ of ferric chloride, ammonium phosphate, and Urea with 40 g L⁻¹ of cow manure). Bar with the same letters is not significantly different ($p \leq 0.05$), Error bars in SEM (standard error mean) N=16.

response to different concentrations of nutrients. Utilizing 15 g L⁻¹ ferric chloride, ammonium phosphate, and urea (group C) and 15 g L⁻¹ ferric chloride, ammonium phosphate, and urea combined with 40 g L⁻¹ cow manure (group D), the maximum cell density reached 86.39×10⁶ cell mL⁻¹ and 85.59×10⁶ cell mL⁻¹, respectively, culture in a mega plastic box (50 L capacity). While increasing SGR of *Nannochloropsis* sp. culture was obtained in group D at 0.308 day⁻¹ after the culture period. Another study conducted bag cultivation (50 L) for microalga *Nannochloropsis oculata* cultured in F/2 medium enriched with different concentrations of nitrogen sources such as sodium nitrate (NaNO₂) and ammonium chloride (NH₄Cl) where the highest number of cells was given as 52×10⁶ cell mL⁻¹ and 49×10⁶ cell mL⁻¹, respectively, at 881 μmol L⁻¹ (Durmaz, 2007). As a result of microalga *N. oculata* culture in a 60 L photobioreactor, harvest cell density in F/2 medium supplemented with 1.76 μmol L⁻¹ of nitrogen was 52.8×10⁶ cell mL⁻¹ (Huang et al., 2013). Low & Toledo (2015) examined that the cell density from 80 L bag culture of microalga *N. oculata* has been harvested at an approximate concentration of 4.55×10⁶ cell mL⁻¹. Additionally, fiberglass reinforced plastic panel photobioreactor for microalga *N. oculata* cultured in F/2 medium obtained the highest density of 245×10⁶ cell mL⁻¹ (Durmaz & Erbil, 2020). It has also been reported that *N. oculata* cultured in 6 L flasks enriched with F/2 medium and supplemented with 100 mg L⁻¹ and 500 mg L⁻¹ concentrations of myo-inositol achieved maximum densities of 101.00×10⁶ cell mL⁻¹ and 91.33×10⁶ cell mL⁻¹, respectively (Erbil & Durmaz, 2020). Other researchers stated that microalga *Porphyridium cruentum* cultured in flat bottom flasks (1 L) enriched with F/2 medium and combined with 0.585×10⁻⁵ M ferric chloride achieved a maximum density of 5.39×10⁶ cell mL⁻¹ (Erbil et al., 2022). Furthermore, it has been demonstrated that high-nutrient manures, such as pig and poultry manures and bi-products of anaerobic digestion, may be viable sources of nutrients for microalgae cultivation (Fenton, 2012; Lu & Xiao, 2022). Our study demonstrated that a combination of cow manure and other inorganic nutrients, such as ferric chloride and ammonium phosphate, significantly increased the microalgae cell density.

CONCLUSION

Nannochloropsis sp. is an important species in aquaculture hatcheries because of its phototrophic nature, enabling a continuous culture of high cell-density organisms. As a result of the present study, the maximum cell density growth of *Nannochloropsis* sp. was significantly increased in both the 15 g L⁻¹ ferric chloride, ammonium phosphate, and urea (group C) and the 15 g L⁻¹ ferric chloride, ammonium phosphate, and urea combined with 40 g L⁻¹ cow manure (group D) at three days of the culture period.

Compliance with Ethical Standards

Authors' Contributions

NBS: Manuscript design, laboratory experiment, and draft checking.

MDH, CTN, RJFR, HAI, JSM, & JHE: Manuscript design and draft checking.

JHS: Statistical analyses, writing, draft checking, reading, and editing.

All authors read and approved the final manuscript.

Conflict of Interest

The authors declare that there is no conflict of interest.

Ethical Approval

For this type of study, formal consent is not required.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

- Barkia, I., Saari, N., & Manning, S. R. (2019). Microalgae for high-value products towards human health and nutrition. *Marine Drugs*, 17(5), 304. <https://doi.org/10.3390/md17050304>
- Benner, P., Meier, L., Pfeffer, A., Krüger, K., Oropeza Vargas, J. E., & Weuster-Botz, D. (2022). Lab-scale photobioreactor systems: Principles, applications, and scalability. *Bioprocess and Biosystems Engineering*, 45(5), 791-813. <https://doi.org/10.1007/s00449-022-02711-1>

- Bogaard, A., Heaton, T. H., Poulton, P., & Merbach, I. (2007). The impact of manuring on nitrogen isotope ratios in cereals: archaeological implications for reconstruction of diet and crop management practices. *Journal of Archaeological Science*, 34(3), 335-343. <https://doi.org/10.1016/j.jas.2006.04.009>
- Chidambara-Murthy, K. N., Vanitha, A., Rajesha, J., Mahadeva-Swamy, M., Sowmya, P. R., & Ravishankar, G. A. (2005). In vivo antioxidant activity of carotenoids from *Dunaliella salina*—a green microalga. *Life Sciences*, 76(12), 1381-1390. <https://doi.org/10.1016/j.lfs.2004.10.015>
- Chu, W. L. (2012). Biotechnological applications of microalgae, *International e-Journal of Science, Medicine & Education*, 6(Suppl 1), S24-S37.
- Ciccione, M. M., Cortese, F., Gesualdo, M., Carbonara, S., Zito, A., Ricci, G., De Pascalis, F., Scicchitano, P., & Riccioni, G. (2013). Dietary intake of carotenoids and their antioxidant and anti-inflammatory effects in cardiovascular care. *Mediators of Inflammation*, 2013, 782137. <https://doi.org/10.1155/2013/782137>
- Dixit, R.B., & Suseela, M. R. (2013). Cyanobacteria: Potential candidates for drug discovery. *Antonie Van Leeuwenhoek*. 103(5), 947-961. <https://doi.org/10.1007/s10482-013-9898-0>
- Duong, V. T., Li, Y., Nowak, E., & Schenk, P. M. (2012). Microalgae isolation and selection for prospective biodiesel production. *Energies*, 5(6), 1835-1849. <https://doi.org/10.3390/en5061835>
- Durmaz, Y., & Erbil, G.Ç. (2017). Effect of light path length of tubes on growth rate of *Nannochloropsis oculata* using industrial scale tubular photobioreactor in the marine hatchery. *Fresenius Environmental Bulletin*, 26(7), 4783-4789.
- Durmaz, Y. (2007). Vitamin E (α -tocopherol) production by the marine microalgae *Nannochloropsis oculata* (Eustigmatophyceae) in nitrogen limitation. *Aquaculture*, 272(1-4), 717-722. <https://doi.org/10.1016/j.aquaculture.2007.07.213>
- Durmaz, Y., & Erbil, G. Ç. (2020). Comparison of industrial-scale tubular photobioreactor to FRP (Fiberglass reinforced plastic) panel photobioreactor on outdoor culture of *Nannochloropsis oculata* in the marine hatchery. *Ege Journal of Fisheries and Aquatic Sciences*, 37(4), 303-308. <https://doi.org/10.12714/egejfas.37.3.13>
- Erbil, G. Ç., & Durmaz, Y. (2020). Effects of myo-inositol concentration on growth and pigments of *Nannochloropsis oculata* culture. *Ege Journal of Fisheries and Aquatic Sciences*, 37(2), 195-199. <https://doi.org/10.12714/egejfas.37.2.11>
- Erbil, G. C., Elp, M., & Durmaz, Y. (2022). Effect of ferric chloride (FeCl₃) concentration on pigment production of *Porphyridium cruentum*. *International Aquatic Research*, 14(2), 127-137. <https://doi.org/10.22034/iar.2022.1950929.1234>
- Fenton, O. (2012). Agricultural nutrient surpluses as potential input sources to grow third generation biomass (microalgae): A review. *Algal Research*, 1(1), 49-56. <https://doi.org/10.1016/j.algal.2012.03.003>
- García-González, A., & Ochoa, J. L. (1999). Anti-inflammatory activity of *Debaryomyces hansenii* Cu, Zn-SOD. *Archives of Medical Research*, 30(1), 69-73. [https://doi.org/10.1016/S0188-0128\(98\)00005-0](https://doi.org/10.1016/S0188-0128(98)00005-0)
- George, B., Pancha, I., Desai, C., Chokshi, K., Paliwal, C., Ghosh, T., & Mishra, S. (2014). Effects of different media composition, light intensity and photoperiod on morphology and physiology of freshwater microalgae *Ankistrodesmus falcatus*—A potential strain for bio-fuel production. *Bioresource Technology*, 171, 367-374. <https://doi.org/10.1016/j.biortech.2014.08.086>
- Giwa, A. (2017). Comparative cradle-to-grave life cycle assessment of biogas production from marine algae and cattle manure biorefineries. *Bioresource Technology*, 244, 1470-1479. <https://doi.org/10.1016/j.biortech.2017.05.143>

- Guzman, S., Gato, A., & Calleja, J. M. (2001). Anti-inflammatory, analgesic and free radical scavenging activities of the marine microalgae *Chlorella stigmatophora* and *Phaeodactylum tricorutum*. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, 15(3), 224-230. <https://doi.org/10.1002/ptr.715>
- Hejazi, M.A., Holwerda, E., & Wijffels, R.H. (2004). Milking microalga *Dunaliella salina* for β -carotene production in two-phase bioreactors. *Biotechnology and Bioengineering*, 85(5), 475-481. <https://doi.org/10.1002/bit.10914>
- Huang, X., Huang, Z., Wen, W., & Yan, J. (2013). Effects of nitrogen supplementation of the culture medium on the growth, total lipid content and fatty acid profiles of three microalgae (*Tetraselmis subcordiformis*, *Nannochloropsis oculata* and *Pavlova viridis*). *Journal of Applied Phycology*, 25, 129-137. <https://doi.org/10.1007/s10811-012-9846-9>
- Huleihel, M., Ishanu, V., Tal, J., & Arad, S. M. (2001). Antiviral effect of red microalgal polysaccharides on *Herpes simplex* and *Varicella zoster* viruses. *Journal of Applied Phycology*, 13(2), 127-134. <https://doi.org/10.1023/A:1011178225912>
- Jjemba, P. K. (2002). The potential impact of veterinary and human therapeutic agents in manure and biosolids on plants grown on arable land: a review. *Agriculture, Ecosystems & Environment*, 93(1-3), 267-278.
- Juneja, A., Ceballos, R. M., & Murthy, G. S. (2013). Effects of environmental factors and nutrient availability on the biochemical composition of algae for biofuels production: A review. *Energies*, 6(9), 4607-4638. <https://doi.org/10.3390/en6094607>
- Lebeau, T., & Robert, J. M. (2003). Diatom cultivation and biotechnologically relevant products. Part I: Cultivation at various scales. *Applied Microbiology and Biotechnology*, 60, 612-623. <https://doi.org/10.1007/s00253-002-1176-4>
- Leu, S., & Boussiba, S. (2014). Advances in the production of high-value products by microalgae. *Industrial Biotechnology*, 10(3), 169-183. <https://doi.org/10.1089/ind.2013.0039>
- Low, C., & Toledo, M. I. (2015). Assessment of the shelf-life of *Nannochloropsis oculata* flocculates stored at different temperatures. *Latin American Journal of Aquatic Research*, 43(2), 315-321. <https://doi.org/10.3856/vol43-issue2-fulltext-7>
- Lu, Q., & Xiao, Y. (2022). From manure to high-value fertilizer: The employment of microalgae as a nutrient carrier for sustainable agriculture. *Algal Research*, 67, 102855. <https://doi.org/10.1016/j.algal.2022.102855>
- Mohan, S. V., Rohit, M. V., Chiranjeevi, P., Chandra, R., & Navaneeth, B. (2015). Heterotrophic microalgae cultivation to synergize biodiesel production with waste remediation: progress and perspectives. *Bioresource Technology*, 184, 169-178. <https://doi.org/10.1016/j.biortech.2014.10.056>
- Mortensen, A. (2006). Carotenoids and other pigments as natural colorants. *Pure and Applied Chemistry*, 78(8), 1477-1491. <https://doi.org/10.1351/pac200678081477>
- Pulz, O., & Gross, W. (2004). Valuable products from biotechnology of microalgae. *Applied Microbiology and Biotechnology*, 65(6), 635-648. <https://doi.org/10.1007/s00253-004-1647-x>
- Skulberg, O. M. (2000). Microalgae as a source of bioactive molecules—experience from cyanophyte research. *Journal of Applied Phycology*, 12(3), 341-348. <https://doi.org/10.1023/A:1008140403621>
- Souza, C. M. M., Bastos, T. S., & dos Santos, M. C. (2021). Microalgae use in animal nutrition. *Research, Society and Development*, 10(16), e53101622986. <https://doi.org/10.33448/rsd-v10i16.22986>